



**VISAGE SURGERY CENTER**  
*Oral & Maxillofacial Surgery*  
*Facial Cosmetic Surgery*  
FAISAL A. QUERESHY, MD, DDS, FACS, INC.  
BOARD CERTIFIED • MEDICAL DIRECTOR

**Post-Operative Instructions**  
**Mentoplasty – Chin Surgery**

**PLEASE READ YOUR INSTRUCTIONS CAREFULLY AND CALL WITH ANY QUESTIONS.**

**POSITION** WHEN RECLINING, ELEVATE YOUR HEAD AND BACK WITH SEVERAL PILLOWS FOR THE FIRST 1-2 DAYS AFTER SURGERY. LIE ON YOUR BACK, RATHER THAN ON YOUR SIDES OR STOMACH.

**BREATHING** IT IS EXTREMELY IMPORTANT THAT YOU TAKE DEEP BREATHS AFTER SURGERY TO PREVENT FROM DEVELOPING PNEUMONIA, WHICH IS A COMMON OCCURRENCE AFTER SURGERY IF BREATHING EXERCISES ARE NOT FOLLOWED PROPERLY. A GOOD EXAMPLE IS TO TAKE 10-20 DEEP BREATHS EVERY HOUR, WHILE YOU ARE AWAKE, AS WELL AS ENCOURAGING YOURSELF TO COUGH.

**ICE** ICE PACKS OR COOL DAMP WASH CLOTS SHOULD BE APPLIED ON OR AROUND THE TREATED AREAS, ESPECIALLY OVER THE EYES AND CHEEKS. THIS WILL LESSEN THE AMOUNT OF SWELLING, BRUISING, AND PAIN. IF NO BANDAGES ARE PRESENT, PLACE A CLOTH BETWEEN THE SKIN AND THE ICE PACK TO PROTECT THE SKIN. USE ICE FOR UP TO 45 MINUTES OUT OF EACH WAKING HOUR FOR THE FIRST 24 HOURS. AN EASY SCHEDULE TO FOLLOW IS 20 MINUTES ON AND 20 MINUTES OFF.

**HEAT** YOU MAY BEGIN HEAT PACKS 12 HOURS AFTER ICE PACKS ARE STOPPED. IF NO BANDAGE IS PRESENT, A MOIST TOWEL SHOULD BE PLACED BETWEEN THE SKIN AND THE HEAT SOURCE. THE MOIST HEAT WILL INCREASE THE CIRCULATION AND HELP THE BODY RID ITSELF OF SWELLING AND BRUISING. DO NOT HEAT CONTINUOUSLY. USE HEAT A MAXIMUM OF 20 MINUTES PER HOUR. THE TEMPERATURE SHOULD BE CLOSELY MONITORED, **NEVER** SET AN ELECTRIC HEATING PAD ABOVE A MEDIUM SETTING, SINCE ANY NUMBNESS IN TREATED AREAS MAKE IT POSSIBLE TO BURN YOUR SKIN WITHOUT KNOWING. TO PREVENT SERIOUS INJURY FROM YOUR ELECTRIC HEATING PAD **BE CERTAIN** IT IS APPROVED FOR USE WITH MOISTURE. HEAT MAY BE USED UNTIL THE SWELLING AND BRUISING HAVE RESOLVED.

**DIET *First Day:*** A CLEAR LIQUID DIET IS RECOMMENDED.

***First Week:*** A LIQUID DIET SHOULD BE FOLLOWED. YOU MAY EAT ANYTHING THAT DOES NOT REQUIRE CHEWING. **ABSOLUTELY NO CHEWING.**

***After One Week:*** SLOWLY RESUME YOUR REGULAR DIET.

**ACTIVITY (I)** DURING YOUR FIRST POST-OPERATIVE DAY, STAY UP AS MUCH AS POSSIBLE. YOU SHOULD SIT, STAND, OR WALK AROUND RATHER THAN REMAIN IN BED. HOWEVER, YOU SHOULD REST WHEN TIRED.

(II) Avoid bending over or lifting more than 5 pounds during the first week.

(III) Take extra precaution to protect your head and neck from bumps, hits or injuries.

**HAIR CARE** YOU MAY WASH YOUR HAIR AFTER SURGERY. A MILD BABY SHAMPOO IS RECOMMENDED. IF YOUR MENTOPLASTY WAS ACCOMPLISHED BY AN INCISION IN THE SKIN UNDER YOUR CHIN, YOU SHOULD AVOID GETTING THE STITCHES OR BANDAGES WET.

**OTHER** DO NOT REMOVE BANDAGES UNLESS OTHERWISE INSTRUCTED.

Following your post-operative appointment when your bandages are removed, you will be instructed how to apply a triangular gauze compression bandage and wrap your head with an ACE elastic bandage to promote rapid healing and decreased swelling. This bandage is to be worn at night and around the house for 3 weeks duration. It may be taken off for work or social events. The bruising will tend to settle in the neck area and can be easily covered with make-up.

Limit talking, as it moves chin muscles.

**IF YOUR MENTOPLASTY WAS ACCOMPLISHED BY AN INCISION INSIDE YOUR MOUTH:** YOUR SURGERY WAS ACCOMPLISHED BY AN INCISION INSIDE YOUR MOUTH. DO NOT DISTURB SUTURES WITH YOUR TONGUE. THE SUTURE CARE ON THE SECOND PAGE IS FOR EXTERNAL SKIN SUTURES ONLY. THE ONLY CARE NECESSARY FOR INTRA-ORAL SUTURES IS THE MOUTH RINSE DESCRIBED BELOW.

*Envision. Entrust. Embark.™*

*Fellow American College of Surgeons • Diplomate American Board of Oral and Maxillofacial Surgery*  
*Board Certified American Board of Cosmetic Surgery • Fellow American Academy of Cosmetic Surgery*



**VISAGE SURGERY CENTER**  
*Oral & Maxillofacial Surgery*  
*Facial Cosmetic Surgery*  
FAISAL A. QUERESHY, MD, DDS, FACS, INC.  
BOARD CERTIFIED • MEDICAL DIRECTOR

Begin oral hygiene three days after surgery. Use the prescribed medicated mouth rinse. Follow directions on bottle. Rinse gently with a capful of this solution 4-5 times a day. DO NOT begin to brush your teeth for five days. A gentle “children’s” toothbrush may be used on your teeth beginning the fifth day after surgery. The oral rinse will help keep the surgical area clean. When you do begin gentle brushing of your teeth, make sure that you do not brush across the incisions with the bristles.

**IF YOUR MENTOPLASTY WAS ACCOMPLISHED BY AN INCISION IN THE SKIN UNDER YOUR CHIN:** NORMAL ORAL HYGIENE ROUTINES MAY BE OBSERVED WITH THE LIMITATION OF NOT OPENING THE MOUTH VERY WIDE DURING BRUSHING. PLEASE SEE INSTRUCTIONS ON THE CARE OF YOUR SKIN STITCHES AT THE END OF THIS SHEET.

**COSMETICS** YOU MAY APPLY COSMETICS TO UNTREATED AREAS FOLLOWING SURGERY. AVOID ANY AREAS WITH SKIN STITCHES. MAKE-UP MAY BE APPLIED AFTER SKIN STITCHES HAVE BEEN REMOVED.

**SPORTS** NO SWIMMING, GYM, OR STRENUOUS ACTIVITIES FOR TWO WEEKS. NO DIVING, BIKING OR SKIING FOR TWO MONTHS. PASSIVE exercise is permitted, like walking. Any other questions should be directed to your surgeon.

**SUN EXPOSURE** PROTECT YOUR FACIAL SKIN FROM EXCESSIVE SUN EXPOSURE AS LONG AS THE TREATED AREA(S) ARE STILL PINK. WHEN THE TREATED AREA(S) ARE NO LONGER PINK, ORDINARY EXPOSURE IS NOT HARMFUL, BUT A SUNSCREEN SHOULD ALWAYS BE USED.

**MEDICATIONS PAIN RELIEVER:** TAKE ONE TABLET WHEN YOU ARRIVE AT HOME. ADDITIONAL TABLETS MAY BE TAKEN EVERY 4-6 HOURS AS NEEDED FOR PAIN RELIEF. **CAUTION:** DO NOT DRIVE OR OPERATE MACHINERY WHILE TAKING PAIN RELIEVERS. TAKE WITH FOOD or liquid to avoid nausea.

**SLEEP MEDICATIONS:** IF PRESCRIBED, TAKE ONE TABLET 30-45 MINUTES PRIOR TO BEDTIME.

**ANTIBIOTICS:** IF PRESCRIBED, TAKE THEM AS INDICATED ON THE BOTTLE AND COMPLETE THE FULL COURSE OF PILLS UNTIL FINISHED.

**ASPIRIN:** AVOID TAKING ASPIRING OR ASPIRING CONTAINING COMPOUNDS DURING YOUR FIRST WEEK AFTER SURGERY.

**ANTI-SWELLING:** (DEXAMETHASONE) IF PRESCRIBED, TAKE ONE TABLET WHEN YOU ARRIVE HOME, AND THEN AGAIN THE FIRST EVENING. THE REMAINING TABLETS SHOULD BE TAKEN OVER THE NEXT TWO DAYS. THESE WILL HELP TO REDUCE THE AMOUNT OF LOCAL swelling.

**CARE OF SKIN STITCHES (TO START AFTER BANDAGE IS REMOVED)**

**SUPPLIES NEEDED:** BACITRACIN OINTMENT  
COTTON TIP APPLICATORS (Q-TIPS, ETC.)  
3% Hydrogen Peroxide (one fresh bottle)  
Normal Saline (one bottle)

**A.** Mix 2 TBSP. PEROXIDE WITH 2 TBSP. NORMA SALINE IN A SMALL CONTAINER. DISCARD EACH TIME, AND DO NOT SAVE MIXTURE.

**B.** Use Q-TIPS AND PEROXIDE SOLUTION TO CLEAN ALL DRY BLOOD AND MATERIAL FORM THE INCISIONS. **DO NOT** LEAVE ANY CRUSTS OR BLOOD ON THE STITCHED AREAS. REPEAT UNTIL SUTURES ARE COMPLETELY DEBRIS FREE AND VISIBLE. REPEAT A MINIMUM OF 4-5 TIMES PER DAY, UNTIL SUTURES ARE REMOVED.

**C.** COVER ALL INCISIONS AND ABRASIONS WITH THIN LAYER OF BACITRACIN OINTMENT – **DO NOT ALLOW ANY AREA** TO DRY OUT OR SCAB OVER. THIS RESULTS IN INCREASED SCARRING.

**D.** DO NOT APPLY ANY BANDAGES OR OTHER MATERIALS TO THE SURGICAL AREA UNLESS OTHERWISE INSTRUCTED.

**PLEASE REPORT ANY OF THE FOLLOWING TO OUR OFFICE:**

**A.** SUDDEN OR EXCESSIVE BLEEDING, SWELLING, OR BRUISING.

**B.** ANY ITCHING, RASH OR REACTION TO MEDICATIONS.

**C.** FEVER, TEMPERATURE OVER 100 DEGREES (TAKEN ORALLY).

**D.** DISCHARGE FROM THE INCISION (OTHER THAN BLOOD).

*Envision. Entrust. Embark.™*

*Fellow American College of Surgeons • Diplomate American Board of Oral and Maxillofacial Surgery  
Board Certified American Board of Cosmetic Surgery • Fellow American Academy of Cosmetic Surgery*



## VISAGE SURGERY CENTER

Oral & Maxillofacial Surgery  
Facial Cosmetic Surgery

---

FAISAL A. QUERESHY, MD, DDS, FACS, INC.  
BOARD CERTIFIED • MEDICAL DIRECTOR

E. ANY INJURY TO THE FACE.

**You May Experience Any or All of the Following:**

**A. SWELLING** – FOLLOWING SURGERY, SWELLING IS A NORMAL BODY RESPONSE. IT USUALLY PEAKS 48 TO 72 HOURS AFTER SURGERY AND SLOWLY RESOLVES DURING THE NEXT 1 TO 2 WEEKS. SLIGHT AMOUNTS OF SWELLING MAY PERSIST FOR UP TO 8 WEEKS.

**B. FACIAL BRUISING** – THIS BECOMES MORE OBVIOUS IN THE SECOND AND THIRD POST OPERATIVE DAY.

**C. NUMBNESS** – PARTS OF YOUR FACE AND CHEEK MAY FEEL WEAK OR NUMB AFTER THE OPERATION. THIS IS USUALLY TEMPORARY.

*FAITHFUL ADHERENCE TO PRE-OPERATIVE AND POST-OPERATIVE INSTRUCTIONS WILL HELP TO MINIMIZE SWELLING, PAIN AND DISCOMFORT. IF YOU DO HAVE ANY PROBLEMS, PLEASE DO NOT HESITATE TO CONTACT ME FOR ASSISTANCE. (330.721.2323).*

**I have received/read and understand the importance of following the above instructions.**

**SIGNED:** \_\_\_\_\_ **DATE:** \_\_\_\_\_

*Envision. Entrust. Embark.™*

*Fellow American College of Surgeons • Diplomate American Board of Oral and Maxillofacial Surgery  
Board Certified American Board of Cosmetic Surgery • Fellow American Academy of Cosmetic Surgery*